

postacią tej alergii. W niniejszej pracy zamieszczamy standardy szczepień dzieci uczulonych na białko jaja kurzego, opracowane na podstawie zaleceń Amerykańskiej Akademii Pediatrii z 2009 r. oraz Brytyjskiego Towarzystwa Alergologii i Immunologii Klinicznej z 2010 r., które są obecnie stosowane w codziennej praktyce 1, 2, 3. Na zachowanie ostrożności w realizacji szczepień przeciwko grypie u dzieci z alergią na białko jaja kurzego wskazują opublikowane niedawno zalecenia Center for Disease Control and Prevention, zamieszczone w MMWR/August 26, 2011/vol. 60/no. 33, 1129-11324.

Abstract:

Each child's immunization may be associated with the risk of adverse reaction. Severe post-vaccination reactions are very rare. Among the large group of children with varying severity of allergy to eggs, the reactions after MMR vaccination can happen, but they are not associated with hypersensitivity to egg protein antigens. The processed protein antigens and their minimum concentration in the MMR vaccine do not cause any allergic reaction. The fixed stereotype thinking of many doctors and parents is disqualification from the MMR vaccination of children allergic to eggs, especially with severe course of allergy. In this paper we present standards of vaccination of children allergic to eggs developed on the basis of the recommendations of the American Academy of Pediatrics in 2009 and the British Society for Allergy and Clinical Immunology in 2010, which are used in everyday practice 1, 2, 3. The caution in the implementation of influenza vaccination in children with allergy to eggs show recently published recommendations of the Center for Disease Control and Prevention published in the MMWR/August 26, 2011/Vol. 60/No. 33, 1129-11324.